

-1-

Date: 1-9-01 Express Mail Label No. EL552288263US

Inventors: Wilhelm Amberg, Teresa Barlozzari, Harald Bernard, Ernst Buschmann,  
Andreas Haupt, Hans-Guenther Hege, Bernd Janssen, Andreas Kling,  
Helmut Lietz, Kurt Ritter, Martina Ullrich, Jürgen Weymann and  
Thomas Zierke

Attorney's Docket No.: 2079.1009-001

## ANTINEOPLASTIC PEPTIDES

### RELATED APPLICATIONS

This application is a continuation of Application No. 09/097,184, filed June 12, 1998, which is a continuation-in-part of International Application No.

- 5 PCT/EP96/05518, filed December 11, 1996, which designated the United States, published in English, which claims priority to U.S. Patent Application No. 08/573,422, filed December 15, 1995, now abandoned. This application also claims benefit of U.S. Provisional Application entitled "Antineoplastic Peptides", which resulted from the conversion of U.S. Serial No. 08/573,422.


- 10 The entire teachings of the above application(s) are incorporated herein by reference.

### FIELD OF THE INVENTION

- The invention described herein provides novel peptides and derivatives thereof which offer potentially improved therapeutic utilities for the treatment of neoplastic
- 15 diseases as compared to dolastatin -10 and -15 (U.S. Patent Nos. 4,879,276 and 4,816,444) and the compounds described in WO 93/23424.

## SUMMARY OF THE INVENTION

Compounds of this invention include novel peptides of the formula I




where

- 5  $R^1$  is hydrogen, methyl, or ethyl;  
 $R^2$  is methyl; or ethyl; or  
 $R^1-N-R^2$  together are a pyrrolidine ring;  
 A is a valyl, isoleucyl, allo-isoleucyl, 2-tert-butylglycyl, 2-ethylglycyl, norleucyl or norvalyl residue;  
 10 B is a N-methyl-valyl, N-methyl-norvalyl, N-methyl-leucyl, N-methyl-isoleucyl, N-methyl-2-tert-butylglycyl, N-methyl-2-ethylglycyl, or N-methyl-norleucyl residue;  
 D is a prolyl, homoprolyl, hydroxyprolyl, or thiazolidine-4-carbonyl residue;  
 15 E is a prolyl, homoprolyl, hydroxyprolyl, thiazolidine-4-carbonyl, trans-4-fluoro-L-prolyl, cis-4-fluoro-L-prolyl, trans-4-chloro-L-prolyl or cis-4-chloro-L-prolyl residue;  
 X is ethyl, propyl, butyl, isopropyl, sec. butyl, tert.-butyl, cyclopropyl, or cyclopentyl;  
 20 G is a L-2-tert.butylglycyl, D-2-terr.butylglycyl, D-valyl, D-isoleucyl, D-leucyl, D-norvalyl, 1-aminopentyl-1-carbonyl, or 2,2-dimethylglycyl residue;  
 s is 0 or 1;  
 K is -NH-C<sub>1-8</sub>-alkyl, -NH-C<sub>3-8</sub>-alkenyl, -NH-C<sub>3-8</sub>-alkinyl, -NH-C<sub>6-8</sub>-cycloalkyl, -NH-C<sub>1-4</sub>-alkene-C<sub>3-8</sub>-cycloalkyl, C<sub>1-4</sub>-alkyl-N-C<sub>1-6</sub>-alkyl, in  
 25 which residues one CH<sub>2</sub> group may be replaced by O or S, one H by phenyl or cyano, or 1, 2 or 3 H by F, except the N-methoxy-N-

-3-

methylamino, N-benzylamino, or N-methyl-N-benzylamino residue, or K  
is



and the salts thereof with physiologically tolerated acids.

## DETAILED DESCRIPTION OF THE INVENTION

In specific embodiments of the compounds of formula I, K may be -NHCH<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>6</sub>CH<sub>3</sub>, -NHCH(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>, -NHCH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>,  
 5 -NHCH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>3</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHCH(CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>3</sub>)C(CH<sub>3</sub>)<sub>3</sub>, -NH-cyclohexyl, -NH-cycloheptyl, -NH-cyclooctyl, -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH(CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>3</sub>)O(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>3</sub>C<sub>6</sub>H<sub>5</sub>, -NHCH(CH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>,  
 10 -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH[CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CN, -NHCH(CH<sub>3</sub>)CH(OH)C<sub>6</sub>H<sub>5</sub>, -NHCH<sub>2</sub>-cyclohexyl, -NHCH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>, -NHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH<sub>2</sub>CF<sub>3</sub>, -NHCH(CH<sub>2</sub>F)<sub>2</sub>, -NHCH<sub>2</sub>CH<sub>2</sub>F, -NHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>2</sub>SCH<sub>3</sub>, -NHCH<sub>2</sub>CHCH<sub>2</sub>, -NH-C(CH<sub>3</sub>)<sub>2</sub>CH=CH<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>C≡CH, -NHC(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>C≡CH,  
 15 -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, -NH(CH<sub>2</sub>CH<sub>2</sub>O)<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -N(OCH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -N(OCH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>, -N(CH<sub>3</sub>)OC<sub>6</sub>H<sub>5</sub>, -NHCH[CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, or the special ring systems mentioned above.

20 In one embodiment of the compounds of formula I described above, s is 0 and E is homopropyl or hydroxypropyl.

Preferred are compounds of the formula I where the substituents R<sup>1</sup>, R<sup>2</sup>, A, B, D, E, X, G and s have the following meanings:

- R<sup>1</sup> hydrogen, methyl, or ethyl, especially methyl;  
 25 R<sup>2</sup> methyl or ethyl, especially methyl;  
 A valyl, isoleucyl, 2-tert-butylglycyl, 2-ethylglycyl, norleucyl or norvalyl, especially valyl, isoleucyl, 2-tert-butylglycyl, 2-ethylglycyl,  
 B N-methyl-valyl, N-methyl-norvalyl, N-methyl-isoleucyl, N-methyl-2-tert-butylglycyl, N-methyl-2-ethylglycyl, or N-methyl-norleucyl, especially N-

methyl-valyl, N-methyl-2-ethylglycyl, N-methyl-norleucyl, N-methyl-isoleucyl, or N-methyl-2-tert.butyl-glycyl;

D prolyl, homoprolyl or thiazolidine-4-carbonyl, especially prolyl or thiazolidine-4-carbonyl;

5 E prolyl, homoprolyl, thiazolidine-4-carbonyl, trans-4-fluoro-L-prolyl, cis-4-fluoro-L-prolyl, trans-4-chloro-L-prolyl or cis-4-chloro-L-prolyl, especially prolyl, trans-4-fluoro-prolyl, cis-4-fluoro-prolyl, trans-4-chloro-prolyl, or cis-4-chloro-prolyl;

X ethyl, propyl, isopropyl, sec.butyl, tert.butyl or cyclo-propyl, especially ethyl, 10 isopropyl, sec.butyl or tert.butyl;


G L-2-tert.butylglycyl, D-2-tert.butylglycyl, D-valyl, D-isoleucyl, D-leucyl or 2,2-dimethylglycyl residue;

s 0 or 1.


Preferred meanings for K are:

15 -NH-C<sub>1-8</sub>-alkyl, -NH-C<sub>6-8</sub>-cycloalkyl, -NH-CH<sub>2</sub>-cyclohexyl, C<sub>1-4</sub>-alkyl-N-C<sub>1-6</sub>-alkyl, in which residues one CH<sub>2</sub> group may be replaced by O, one H by phenyl or 1 or 2 H by F, except the N-methoxy-N-methylamino, N-benzylamino, or N-methyl-N-benzylamino residue, or K is


20



25




-6-



More preferred K is


- NHCH<sub>3</sub>, -NHCH<sub>2</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>4</sub>CH<sub>3</sub>,  
 10 -NH(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>6</sub>CH<sub>3</sub>, -NH(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>, -NHCH(CH<sub>3</sub>)<sub>2</sub>,  
 -NHCH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>3</sub>,  
 -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHCH(CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>,  
 -NHCH(CH<sub>3</sub>)C(CH<sub>3</sub>)<sub>3</sub>, -NH-cyclohexyl, -NH-cycloheptyl, -NH-cyclooctyl,  
 -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH(CH<sub>3</sub>)<sub>2</sub>,  
 15 -N(OCH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>3</sub>)OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>3</sub>C<sub>6</sub>H<sub>5</sub>,  
 -NHCH(CH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>,  
 -NHCH(CH<sub>3</sub>)CH(OH)C<sub>6</sub>H<sub>5</sub>, -NHCH<sub>2</sub>-cyclohexyl, -N(CH<sub>3</sub>)<sub>2</sub>, -N(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>,  
 -N(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>F)<sub>2</sub>, -NHC(CH<sub>3</sub>)CH = CH<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CN,  
 -NHC(CH<sub>3</sub>)<sub>2</sub>C ≡ CH, -NHC(CH<sub>3</sub>)<sub>2</sub>CONH<sub>2</sub>, -NHCH[CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>,  
 20 -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,




In one embodiment of the preferred compounds of formula I described above, s is 0 and E is homoprolyl or hydroxyprolyl.

Especially preferred are compounds of the formula I where

- R<sup>1</sup> and R<sup>2</sup> are methyl,
- 5 A is a valyl, isoleucyl, 2-tert.-butylglycyl residue
- B is a N-methylvalyl, N-methyl-isoleucyl, N-methyl-2-tert.-butylglycyl residue,
- D is a prolyl or thiazolidine-4-carbonyl residue,
- E is a prolyl, cis-4-fluoro-L-prolyl, or cis-4-chloro-L-prolyl residue,
- X is a isopropyl, sec.-butyl, or tert.-butyl residue,
- 10 s is 0, and
- K is
- NHCH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>,  
 -NHC(CH<sub>3</sub>)<sub>3</sub>, -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHCH(CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>,  
 -NHCH(CH<sub>2</sub>CH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>, -NHCH(CH<sub>3</sub>)C(CH<sub>3</sub>)<sub>3</sub>, -NH-cycloheptyl, -NH-cyclooctyl,
- 15 -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)OCH(CH<sub>3</sub>)<sub>2</sub>, -N(OCH<sub>3</sub>)CH(CH<sub>3</sub>)<sub>2</sub>,  
 -N(CH<sub>3</sub>)OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NH(CH<sub>2</sub>)<sub>3</sub>C<sub>6</sub>H<sub>5</sub>, -NHCH(CH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>,  
 -NHC(CH<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>,  
 -NHCH(CH<sub>3</sub>)CH(OH)C<sub>6</sub>H<sub>5</sub>, -NHCH(CH<sub>2</sub>F)<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH,  
 -NH(CH<sub>2</sub>CH<sub>2</sub>O)<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH=CH<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>,
- 20 -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
 -NHC(CH<sub>3</sub>)<sub>2</sub>CN, -NHC(CH<sub>3</sub>)<sub>2</sub>C≡CH, -NHCH[CH(CH<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, -NHC(CH<sub>3</sub>)<sub>2</sub>CONH<sub>2</sub>,  
 -NHC(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -N(OCH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>, -N(OCH<sub>3</sub>)CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>,





This invention also provides methods for preparing the compounds of formula I, pharmaceutical compositions containing such compounds together with a  
 10 pharmaceutically acceptable carrier and methods for using same for treating cancer in mammals.

The new compounds may be present as salts with physiologically tolerated acids such as: hydrochloric acid, citric acid, tartaric acid, lactic acid, phosphoric acid, methanesulfonic acid, acetic acid, formic acid, maleic acid, fumaric acid, malic acid,  
 15 succinic acid, malonic acid, sulfuric acid, L-glutamic acid, L-aspartic acid, pyruvic acid, mucic acid, benzoic acid, glucuronic acid, oxalic acid, ascorbic acid and acetyl glycine.

The novel compounds can be prepared by known methods of peptide chemistry. Thus, the peptides can be assembled sequentially from amino acids or by linking suitable small peptide fragments. In the sequential assemblage, starting at the C  
 20 terminus the peptide chain is extended stepwise by one amino acid each time. In fragment coupling it is possible to link together fragments of different lengths, and the fragments in turn can be obtained by sequential assemblage from amino acids or themselves by fragment-coupling.

Both in the sequential assemblage and in the fragment coupling it is necessary to  
 25 link the units by forming an amide linkage. Enzymatic and chemical methods are suitable for this.

Chemical methods for forming the amide linkage are described in detail by Mueller, Methoden der organischen Chemie Vol. XV/2, pp 1 to 264, Thieme Verlag, Stuttgart, 1974; Stewart, Young, Solid Phase Peptide Synthesis, pp 31 to 34, 71 to 82,



- Pierce Chemical Company, Rockford, 1984; Bodanszky, Klausner, Ondetti, Peptide Synthesis, pp 85 to 128, John Wiley & Sons, New York, 1976; The Practice of Peptide Synthesis, M. Bodanszky, A. Bodanszky, Springer-Verlag, 1994, and other standard works on peptide chemistry. Particular preference is given to the azide method, the
- 5 symmetric and mixed anhydride method, in situ generated or preformed active esters, the use of urethane protected N-carboxy anhydrides of amino acids and the formation of the amide linkage using coupling reagents, especially dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), 1-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), pivaloylchloride, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
  - 10 hydrochloride (EDCI), n-propanephosphonic anhydride (PPA), N,N-bis(2-oxo-3-oxazolodiny)-amidophosphoryl chloride (BOP-C1), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBrop), diphenylphosphoryl azide (DPPA), Castro's reagent (BOP, PyBop), O-benzotriazolyl-N,N,N',N'-tetramethyluronium salts (HBTU), O-azabenzotriazolyl-N,N,N',N'-tetramethyluronium salts (HATU),
  - 15 diethylphosphoryl cyanide (DEPCN), 2,5-diphenyl-2,3-dihydro-3-oxo-4-hydroxythiophene dioxide (Steglich's reagent; HOTDO) and 1,1'-carbonyldiimidazole (CDI). The coupling reagents can be employed alone or in combination with additives such as N,N-dimethyl-4-aminopyridine (DMAP), N-hydroxy-benzotriazole (HOBt), N-hydroxybenzotriazine (HOObt), Azabenzotriazole, N-hydroxysuccinimide (HOSu) or
  - 20 2-hydroxypyridine.

Whereas it is normally possible to dispense with protective groups in enzymatic peptide synthesis, reversible protection of reactive groups not involved in formation of the amide linkage is necessary for both reactants in chemical synthesis. Three conventional protective group techniques are preferred for the chemical peptide

- 25 synthesis: the benzyloxycarbonyl (Z), the t-butoxycarbonyl (Boc) and the 9-fluorenylmethoxycarbonyl (Fmoc) techniques.

Identified in each case is the protective group on the alpha-amino group of the chain-extending unit. A detailed review of amino-acid protective groups is given by Mueller, Methoden der organischen Chemie vol. XV/1, pp 20 to 906, Thieme Verlag,

Stuttgart, 1974. The units employed for assembling the peptide chain can be reacted in solution, in suspension or by a method similar to that described by Merrifield in J. Amer. Chem. Soc. 85 (1963) 2149.

Suitable for peptide synthesis in solution are all solvents which are inert under  
5 the reaction conditions, especially water, N,N-dimethylformamide (DMF), dimethyl sulfoxide (DMSO), acetonitrile, dichloromethane (DCM), ethyl acetate, 1,4-dioxane, tetrahydrofuran (THF), N-methyl-2-pyrrolidone (NMP) and mixtures of the said solvents.

Peptide synthesis on the polymeric support can be carried out in all inert organic  
10 solvents in which the amino-acid derivatives used are soluble. However, preferred solvents additionally have resin-swelling properties, such as DMF, DCM, NMP, acetonitrile and DMSO, and mixtures of these solvents. After synthesis is complete, the peptide is cleaved off the polymeric support. The conditions under which cleavage off the various resin types is possible are disclosed in the literature. The cleavage reactions  
15 most commonly used are acid- and palladium-catalyzed, especially cleavage in liquid anhydrous hydrogen fluoride, in anhydrous trifluoromethanesulfonic acid, in dilute or concentrated trifluoroacetic acid, palladium-catalyzed cleavage in THF or THF-DCM mixtures in the presence of a weak base such as morpholine or cleavage in acetic acid/dichloromethane/ trifluoroethanol mixtures. Depending on the chosen protective  
20 groups, these may be retained or likewise cleaved off under the cleavage conditions.

Partial deprotection of the peptide may also be worthwhile when certain derivatization reactions are to be carried out.

Peptides dialkylated at the N-terminus can be prepared either by coupling on the appropriate N,N-di-alkylamino acids in solution or on the polymeric support, by  
25 reductive alkylation of the resin-bound peptide in DMF/1% acetic acid with  $\text{NaCNBH}_3$  and the appropriate aldehydes, by hydrogenation of the peptide in solution in the presence of aldehyde or ketone and Pd/C.

The various non-naturally occurring amino acids as well as the various non-amino acid moieties disclosed herein may be obtained from commercial sources or

synthesized from commercially-available materials using methods known in the art. For example, amino acids building blocks with  $R^1$  and  $R^2$  moieties can be prepared according to E. Wuensch, Houben Weyl, Meth. d. Org. Chemie, Bd. XV, 1, p. 306 following, Thieme Verlag Stuttgart 1974 and Literature cited therein.

- 5           The compounds of this invention may be used to inhibit or otherwise treat solid tumors (e.g. tumors of the lung, breast, colon, prostate, bladder, rectum, or endometrial tumors) or hematological malignancies (e.g. leukemias, lymphomas) by administration of the compound to the mammal.

          It is a special advantage of the new compounds that they are very resistant to  
10 enzymatic degradation and can also be administered orally.

          Administration may be by any of the means which are conventional for pharmaceutical, preferably oncological, agents, including oral and parenteral means such as subcutaneously, intravenously, intramuscularly and intraperitoneally.

- The compounds may be administered alone or in the form of pharmaceutical  
15 compositions containing a compound of formula I together with a pharmaceutically accepted carrier appropriate for the desired route of administration. Such pharmaceutical compositions may be combination products, i.e., may also contain other therapeutically active ingredients.

- The dosage to be administered to the mammal will contain an effective tumor-  
20 inhibiting amount of active ingredient which will depend upon conventional factors including the biological activity of the particular compound employed; the means of administration; the age, health and body weight of the recipient; the nature and extent of the symptoms; the frequency of treatment; the administration of other therapies; and the effect desired. A typical daily dose will be about 0.05 to 50 milligrams per kilogram of  
25 body weight on oral administration and about 0.01 to 20 milligrams per kilogram of body weight on parenteral administration.

          The novel compounds can be administered in conventional solid or liquid pharmaceutical administration forms, e.g. uncoated or (film-)coated tablets, capsules, powders, granules, suppositories or solutions. These are produced in a conventional

manner. The active substances can for this purpose be processed with conventional pharmaceutical aids such as tablet binders, fillers preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, sustained release compositions, antioxidants and/or propellant gases (cf. H. Sucker et al.:

- 5 Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1978). The administration forms obtained in this way normally contain 1-90% by weight of the active substance.

The following examples are intended to illustrate the invention. The proteinogenous amino acids are abbreviated in the examples using the known three-letter code. Other abbreviations used: Me<sub>2</sub>Val = N,N-dimethylvaline, MeVal =

- 10 N-methylvaline.

## EXAMPLES

### A. General procedures

- I. The peptides claimed in claim 1 are either synthesized by classical solution  
15 synthesis using standard Z- and Boc-methodology as described above or by standard methods of solid-phase synthesis using Boc and Fmoc protective group techniques.

- In the case of solid phase synthesis, the N,N-dialkylpenta- or hexapeptide acids  
20 are liberated from the solid support and further coupled with the corresponding C-terminal amines in solution. BOP-C1 and PyBrop were used as reagents for coupling of the amino acid following the N-methylamino acids. The reaction times were correspondingly increased. For reductive alkylation of the N-terminus, the peptide-resin was deprotected at the N terminus and then reacted  
25 with a 3-fold molar excess of aldehyde or ketone in DMF/1% acetic acid with addition of 3 equivalents of NaCNBH<sub>3</sub>. After the reaction was complete (negative Kaisertest) the resin was washed several times with water, isopropanol, DMF and dichloromethane.

In solution synthesis, the use of either Boc-protected amino acid NCAs (N-tert.-butyloxycarbonyl-amino acid-N-carboxy-anhydrides), Z-protected amino acid NCAs (N-benzyloxycarbonyl-amino acid-N-carboxy-anhydrides), or the use of pivaloylchloride as condensing agent respectively is most advantageous for coupling of the amino acid following the N-methylamino acids. Reductive alkylation of the N terminus can e.g. be achieved by reaction of the N-terminally deprotected peptides or amino acids with the corresponding aldehydes or ketones using  $\text{NaCNBH}_3$  or hydrogen, Pd/C.

## 10 II. Purification and characterization of the peptides

Purification was carried out by gel chromatography (SEPHADEX G-10, G-15/10% HOAc, SEPHADEX LH20/MeOH), medium pressure chromatography (stationary phase: HD-SIL C-18, 20-45 mikron, 100 Angstrom; mobile phase: gradient with A = 0.1% TFA/MeOH, B = 0.1% TFA/water), or preparative HPLC (stationary phase: Waters Delta-Pak C-18, 15 mikron, 100 Angstrom; mobile phase: gradient with A = 0.1% TFA/MeOH, 3 = 0.1% TFA/water).

The purity of the resulting products was determined by analytical HPLC (stationary phase: 100 2.1 mm VYDAC C-18, 5  $\mu$ , 300 A; mobile phase: acetonitrile-water gradient, buffered with 0.1% TFA, 40°C).

Characterization was by amino-acid analysis and fast atom bombardment mass spectroscopy.

## B. Specific procedures

## EXAMPLE 1 (SEQ ID NO: 1)

 $\text{Me}_2\text{Val}-\text{Val}-\text{MeVal}-\text{Pro}-\text{Pro}-\text{NHCH}(\text{CH}_3)_2$ 

## a) Z-MeVal-Pro-OMe

5 66.25 g (250 mmol) Z-MeVal-OH were dissolved in 250 ml dry dichloromethane. After addition of 36.41 ml (262.5 mmol) triethylamine, the reaction mixture was cooled to  $-25^\circ\text{C}$  and 32.27 ml (262.5 mmol) pivaloyl chloride were added. After stirring for 2,5 h, 41.89 g (250 mmol) H-Pro-OMe x Chl in 250 ml dichloromethane, neutralized with 36.41 ml (262.5 mmol)

10 triethylamine at  $0^\circ\text{C}$ , were added to the reaction mixture. Stirring continued for 2 h at  $-25^\circ\text{C}$  and overnight at room temperature. The reaction mixture was diluted with dichloromethane and thoroughly washed with saturated aqueous  $\text{NaHCO}_3$  solution (3x), water (1x), 5 % citric acid (3x) and saturated NaCl solution. The organic phase was dried over sodium sulfate and evaporated to

15 dryness. The residue (91.24 g) was stirred with petroleum ether overnight and filtered. 62.3 g of product were obtained.

## b) H-MeVal-Pro-OMe

48.9 g (130 mmol) Z-MeVal-Pro-OMe were dissolved in 490 ml methanol.

20 After addition of 10.9 ml (130 mmol) concentrated hydrochloric acid and 2.32 g 10 % Palladium/charcoal, the reaction mixture was hydrogenated. Filtration and evaporation to dryness yielded 36.32 g of the product.

## c) Z-Val-MeVal-Pro-OMe

25 18.1 g (65 mmol) H-MeVal-Pro-OMe, 21.6 g (78 mmol) Z-Val-N-carboxyanhydride and 22.8 ml (130 mmol) diisopropylethylamine were stirred in 110 ml DMF at  $40^\circ\text{C}$  for 2 d. After evaporation of DMF, dichloromethane was added and the organic phase washed with saturated aqueous  $\text{NaHCO}_3$  solution (3x), water (1x), 4 % citric acid (3X) and saturated NaCl solution. The organic

phase was dried over sodium sulfate and evaporated to dryness. The product (29.3 g) was obtained as a viscous oil.

d) H-Val-MeVal-Pro-OMe

5 29.3 g (61.6 mmol) of Z-Val-MeVal-Pro-OMe were dissolved in 230 ml methanol. After addition of 1.15 g 10% Palladium/charcoal, the reaction mixture was hydrogenated. Filtration and evaporation to dryness yielded 21.96 g of the product.

10 e) Z-Val-Val-MeVal-Pro-OMe

15 15.29 g (61 mmol) Z-Val-OH and 21.96 g (61 mmol) H-Val-MeVal-Pro-OMe were dissolved in 610 ml dichloromethane and cooled to 0°C. After addition of 8.16 ml (73.2 mmol) N-Methylmorpholine, 2.77 g (20.3 mmol) HOBt and 11.73 g (61 mmol) EDCI, the reaction mixture was stirred overnight at room temperature, diluted with dichloromethane and thoroughly washed with saturated aqueous NaHCO<sub>3</sub> solution (3x), water (1x), 5 % citric acid (3X) and saturated NaCl solution. The organic phase was dried over sodium sulfate and evaporated to dryness to yield 31.96 g of the product.

20 f) Z-Val-Val-MeVal-Pro-OH

25 31.96 g (57 mmol) Z-Val-Val-MeVal-Pro-OMe were dissolved in 250 ml methanol. 102.6 ml of a 1 N LiOH solution was added and the mixture stirred overnight at room temperature. After addition of 500 ml water, the aqueous phase was washed three times with ethyl acetate, adjusted to pH 2 at 0° C and extracted three times with ethyl acetate. The organic phase was dried over sodium sulfate and evaporated to dryness yielding 30.62 g of the desired product as a white solid.

g) Z-Val-Val-MeVal-Pro-Pro-NHCH(CH<sub>3</sub>)<sub>2</sub>

2 g (3.35 mmol) Z-Val-Val-MeVal-Pro-OH and 0.664 g (3.35 mmol) H-Pro-NHCH(CH<sub>3</sub>)<sub>2</sub> were dissolved in 34 ml of dry dichloromethane. After cooling to 0° C, 1.35 ml (12.1 mmol) N-methylmorpholine, 0.114 g (0.84 mmol) HOBt and 0.645 g (3.35 mmol) EDCI were added and the reaction mixture stirred overnight at room temperature. 80 ml dichloromethane were added and the organic phase thoroughly washed with saturated aqueous NaHCO<sub>3</sub> solution (3x), water (1x), 5 % citric acid (3x) and saturated NaCl solution (1x). The organic phase was dried over sodium sulfate and evaporated to dryness to yield 1.96 g of the product which was used in the next reaction without further purification.

h) Me<sub>2</sub>Val-Val-MeVal-Pro-Pro-NHCH(CH<sub>3</sub>)<sub>2</sub>

1.96 g Z-Val-Val-MeVal-Pro-Pro-NHCH(CH<sub>3</sub>)<sub>2</sub> were dissolved in 11 ml methanol. 0.054 g 10 % Pd/C were added under nitrogen atmosphere and the reaction mixture hydrogenated at room temperature for 4 h. After addition of 0.86 ml (11.24 mmol) of a 37 % aqueous formaldehyde solution and 0.281 g 10 % Pd/C, hydrogenation was continued for 5 h. Filtration and evaporation of the solvent gave rise to 2.77 g of crude product. Further purification was achieved by dissolving the peptide in water, adjusting the pH to 2 and extracting the aqueous phase three times with ethyl acetate. The aqueous phase was then adjusted to pH 8-9 and extracted four times with dichloromethane. The organic phase was dried over sodium sulfate to yield 1.37 g of purified product as a white foam. The compound was further purified using medium pressure liquid chromatography (10 - 50 % A in 10 min.; 50 - 90 % A in 320 min.). Fractions containing the product were combined, lyophilized, redissolved in water and the pH adjusted to 9 with 1 N LiOH. After extraction with dichloromethane, the organic phase was dried over sodium sulfate and evaporated to dryness. Lyophilization led to 500 mg of pure product, which was characterized by fast atom bombardment mass spectrometry ([M+H]<sup>+</sup> = 593).



## EXAMPLE 2 (SEQ ID NO: 1)

Me<sub>2</sub>Val-Val-MeVal-Pro-Pro-NHC(CH<sub>3</sub>)<sub>3</sub>

i) Z-Val-Val-MeVal-Pro-Pro-NHC(CH<sub>3</sub>)<sub>3</sub>

2 g (3.35 mmol) Z-Val-Val-MeVal-Pro-OH and 0.692 g (3.35 mmol) H-Pro-NHC(CH<sub>3</sub>)<sub>3</sub> were dissolved in 34 ml of dry dichloromethane. After cooling to 0°C, 1.35 ml (12.1 mmol) N-methylmorpholine, 0.114 g (0.84 mmol) HOBt and 0.645 g (3.35 mmol) EDCI were added and the reaction mixture stirred overnight at room temperature. 80 ml dichloromethane were added and the organic phase thoroughly washed with saturated aqueous NaHCO<sub>3</sub> solution (3x), water (1x), 5 % citric acid (3x) and saturated NaCl solution (1x). The organic phase was dried over sodium sulfate and evaporated to dryness to yield 1.8 g of the product which was used in the next reaction without further purification.

k) Me<sub>2</sub>Val-Val-MeVal-Pro-Pro-NHC(CH<sub>3</sub>)<sub>3</sub>

1.8 g Z-Val-Val-MeVal-Pro-Pro-NHC(CH<sub>3</sub>)<sub>3</sub> were dissolved in 10 ml methanol. 0.049 g 10 % Pd/C were added under nitrogen atmosphere and the reaction mixture hydrogenated at room temperature for 4 h. After addition of 0.86 ml (11.24 mmol) of a 37 % aqueous formaldehyde solution and 0.252 g 10 % Pd/C, hydrogenation was continued for 5 h. Filtration and evaporation of the solvent gave rise to 1.82 g of crude product. The compound was further purified using medium pressure liquid chromatography 910 - 50 % A in 10 min.; 50 - 90 % A in 320 min.). Fractions containing the product were combined, lyophilized, redissolved in water and the pH adjusted to 9 with 1 N LiOH. After extraction with dichloromethane, the organic phase was dried over sodium sulfate and evaporated to dryness. Lyophilization led to 547 mg of pure product, which was characterized by fast atom bombardment mass spectrometry ([M+H]<sup>+</sup> = 607).

The following compounds were prepared or can be prepared according to examples 1 and 2:

- |    |     |                     |
|----|-----|---------------------|
|    | 3.  | Xaa Val Xab Pro Xac |
| 5  | 4.  | Xaa Val Xab Pro Xad |
|    | 5.  | Xaa Val Xab Pro Xae |
|    | 6.  | Xaa Val Xab Pro Xaf |
|    | 7.  | Xaa Val Xab Pro Xag |
|    | 8.  | Xaa Val Xab Pro Xah |
|    | 9.  | Xaa Val Xab Pro Xai |
|    | 10. | Xaa Val Xab Pro Xak |
| 10 | 11. | Xaa Val Xab Pro Xal |
|    | 12. | Xaa Val Xab Pro Xam |
|    | 13. | Xaa Val Xab Pro Xan |
|    | 14. | Xaa Val Xab Pro Xao |
|    | 15. | Xaa Val Xab Pro Xap |
|    | 16. | Xaa Val Xab Pro Xaq |
|    | 17. | Xaa Val Xab Pro Xar |
| 15 | 18. | Xaa Val Xab Pro Xas |
|    | 19. | Xaa Val Xab Pro Xat |
|    | 20. | Xaa Val Xab Pro Xau |
|    | 21. | Xaa Val Xab Pro Xav |
|    | 22. | Xaa Val Xab Pro Xaw |
|    | 23. | Xaa Val Xab Pro Xax |
|    | 24. | Xdd Val Xab Pro Xay |
| 20 | 25. | Xaa Val Xab Pro Xaz |
|    | 26. | Xaa Val Xab Pro Xba |
|    | 27. | Xaa Val Xab Pro Xbb |
|    | 28. | Xaa Val Xbc Pro Xay |
|    | 29. | Xaa Val Xab Pro Xbd |
|    | 30. | Xaa Val Xab Pro Xbe |
|    | 31. | Xaa Val Xab Pro Xbf |
| 25 | 32. | Xaa Val Xab Pro Xbg |
|    | 33. | Xaa Val Xab Pro Xbh |
|    | 34. | Xaa Val Xab Pro Xbi |
|    | 35. | Xaa Val Xab Pro Xbk |
|    | 36. | Xaa Val Xab Pro Xbl |
|    | 37. | Xaa Val Xab Pro Xbm |
|    | 38. | Xaa Val Xab Pro Xbn |

	39.	Xaa Val Xab Pro Xbo
	40.	Xaa Val Xab Pro Xbp
	41.	Xaa Val Xab Pro Xbq
	42.	Xaa Val Xab Pro Xbr
	43.	Xaa Val Xab Pro Xbs
5	44.	Xaa Val Xab Pro Xbt
	45.	Xaa Val Xab Pro Xbu
	46.	Xaa Val Xab Pro Xbv
	47.	Xaa Val Xab Pro Xbw
	48.	Xaa Val Xab Pro Xbx
	49.	Xaa Val Xab Pro Xby
	50.	Xaa Val Xab Pro Xbz
10	51.	Xaa Val Xab Pro Xca
	52.	Xaa Val Xab Pro Xcb
	53.	Xaa Val Xab Pro Xcc
	54.	Xaa Val Xab Pro Xcd
	55.	Xaa Val Xab Pro Xce
	56.	Xaa Val Xab Pro Xcf
	57.	Xaa Xdf Xab Pro Xay
15	58.	Xaa Val Xab Pro Xch
	59.	Xaa Val Xab Pro Xci
	60.	Xaa Val Xab Pro Xck
	61.	Xaa Val Xab Pro Xcl
	62.	Xaa Val Xab Pro Xcm
	63.	Xaa Val Xab Pro Xcn
	64.	Xaa Val Xab Pro Xco
20	65.	Xaa Val Xab Pro Xcp
	66.	Xaa Val Xab Pro Xcq
	67.	Xaa Val Xab Pro Xcr
	68.	Xaa Val Xab Pro Xcs
	69.	Xaa Val Xab Pro Xct
	70.	Xaa Val Xab Pro Xcu
	71.	Xcw Val Xab Pro Xcv
25	72.	Xcx Val Xab Pro Xcv
	73.	Xaa Val Xab Pro Pro Xcy
	74.	Xaa Val Xab Pro Pro Xcz

- 5 75. Xaa Val Xda Pro Xcv  
76. Xaa Xdb Xab Pro Xcv  
77. Xdc Val Xab Pro Xcv  
78. Xaa Ile Xab Pro Xcv  
79. Xdd Val Xab Pro Xcv  
80. Xde Val Xab Pro Xcv  
81. Xaa Xdf Xab Pro Xcv  
82. Xaa Val Xab Pro Xcg  
83. Xaa Val Xab Pro Pro Xdg  
84. Xaa Val Xab Pro Pro Xdh  
85. Xaa Val Xab Pro Pro Xdi  
86. Xaa Val Xab Pro Pro Xdk
- 10 87. Xaa Val Xdl Pro Xcv  
88. Xde Val Xab Pro Xay  
89. Xaa Val Xdl Pro Xay  
90. Xaa Val Xab Pro Xdm  
91. Xaa Val Xab Pro Xdn  
92. Xaa Val Xab Pro Xdo  
93. Xaa Val Xab Pro Xdp  
94. Xaa Val Xab Pro Xdq
- 15 95. Xaa Val Xab Pro Pro Xdr  
96. Xaa Val Xab Pro Xds  
97. Xaa Val Xbc Pro Xcv  
98. Xaa Ile Xab Pro Xay  
99. Xcw Val Xab Pro Xay  
100. Xaa Val Xbc Pro Xal
- 20 101. Xaa Val Xdl Pro Xal  
102. Xaa Xdf Xab Pro Xal  
103. Xaa Ile Xab Pro Xal  
104. Xdd Val Xab Pro Xal  
105. Xde Val Xab Pro Xal  
106. Xcx Val Xab Pro Xcy  
107. Xcw Val Xab Pro Xal
- 25 108. Xcx Val Xab Pro Xal  
109. Xcw Val Xab Pro Xav  
110. Xcx Val Xab Pro Xav  
111. Xcw Val Xab Pro Xaw  
112. Xcx Val Xab Pro Xaw  
113. Xab Val Xab Pro Xay  
114. Xab Val Xab Pro Xcv

5 115. Xab Val Xab Pro Xal  
116. Xab Val Xab Pro Xam  
117. Xab Val Xab Pro Xan  
118. Xab Val Xab Pro Xao  
119. Xab Val Xab Pro Xav  
120. Xab Val Xab Pro Xaw  
121. Xab Val Xab Pro Xaz  
122. Xab Val Xab Pro Xau  
123. Xab Val Xab Pro Xbf  
124. Xab Val Xab Pro Xbm  
125. Xab Val Xab Pro Xbn  
10 126. Xab Val Xab Pro Xbo  
127. Xab Val Xab Pro Xch  
128. Xaa Val Xab Pro Xdt  
129. Xaa Val Xab Pro Xdu  
130. Xaa Val Xab Pro Xdv  
131. Xaa Val Xab Pro Xdw  
132. Xaa Val Xab Pro Xdx  
133. Xaa Val Xab Pro Xdy  
15 134. Xaa Val Xab Pro Xdz  
135. Xaa Val Xab Pro Xea  
136. Xaa Val Xab Pro Xeb  
137. Xaa Val Xab Pro Xec  
138. Xaa Val Xab Pro Xed  
139. Xaa Val Xab Pro Xef  
140. Xaa Val Xab Pro Xeg  
20 141. Xaa Val Xab Pro Xeh  
142. Xaa Val Xab Pro Xei  
143. Xaa Val Xab Pro Xek  
144. Xaa Val Xab Pro Xel  
145. Xaa Val Xab Pro Xem  
146. Xaa Val Xab Pro Xen  
147. Xaa Val Xab Pro Xeo  
25 148. Xaa Val Xab Pro Xep  
149. Xaa Val Xab Pro Xeq  
150. Xaa Val Xab Pro Xer  
151. Xaa Val Xab Pro Xcg

Examples for the MS-characterization of the synthesized novel compounds are given in the following table.

	EXAMPLE	Fast atom bombardment MS analysis.
	[No.]	[Mol.-Weight (measured)]
5	3.	565
	4.	579
	5.	593
	6.	607
	7.	621
10	8.	635
	11.	607
	12.	607
	13.	621
	14.	649
15	15.	635
	16.	635
	17.	635
	18.	635
	19.	621
20	20.	621
	21.	635
	22.	635
	25.	633
	26.	647
25	27.	661
	31.	623
	32.	671
	33.	667
	34.	631
	35.	655
	36.	655
	37.	669

-23-

## EXAMPLE

Fast atom bombardment MS analysis.

5	38.	621
	39.	635
	41.	649
	42.	621
	43.	633
	44.	667
	45.	607
	46.	647
	47.	668
	48.	655
10	49.	669
	50.	685
	51.	629
	52.	625
	53.	721
15	55.	579
	58.	623
	61.	597
	62.	621
	63.	609
20	64.	625
	65.	635
	66.	591
	67.	715
	68.	685
25	69.	685
	70.	591
	71.	607
	72.	621
	74.	706
	75.	579
	76.	579
	77.	579

-24-

EXAMPLE		Fast atom bombardment MS analysis.
5	73.	607
	79.	607
	80.	607
	81.	607
	82.	637
10	83.	692
	84.	706
	85.	706
	86.	706
	87.	607
15	90.	635
	92.	659
	93.	617
	94.	636
	95.	678
20	128.	671
	131.	625
	139	625
25	151.	637

Table I - Sequence Identification of Compounds Prepared According to Examples 1 and 2

Compound Number(s)	Sequence ID Number
1-56, 58-72, 75, 77, 79, 80, 82, 87-94, 96, 97, 99-101, 104-151	1
73, 74, 83-86, 95,	2
57, 76, 81, 102	3
78, 98, 103	4


The symbols Xaa in the summary have the following meanings:

Xaa: N,N-Dimethylvaline  
 Xab: N-Methylvaline




Xac:

5




Xad:

10




Xae:

15




Xaf:

20




Xag:

25




Xah:




5

Xai:




10

Xak:




15

Xal:




20

Xam:




Xan:



25


Xao:

5




Xap:

10




Xaq:

15




Xar:

20




Xas:

25




Xat:

5




Xau:

10




Xav:

15




Xaw:

20




Xax:

25




Xay:

5




Xaz:

10




Xba:

15



Xbb:

20




Xbc:

N-Methyl-isoleucine


25

Xbd:




-30-

Xbe:




5

Xbf:




10

Xbg:




15

Xbh:




20

Xbi:




25

Xbk:




-31-

Xb1:




5

Xbm:




10

Xbn:




15

Xbo:



20


Xbp:



25


Xbq:

5




Xbr:

10




Xbs:

15




Xbt:

20




Xbu:

25




Xbv:






Xbw

5




Xbx:

10




Xby:

15




Xbz:

20




Xca:

25



-34-

Xcb:




5

Xcc:


Proline adamantyl (1) amide

Xcd:




10

Xce:




15

Xcf:




20

Xcg:




25

Xch:




5

Xci:




10

Xck:




15

Xcl:



20


Xcm:



25


-36-

Xcn:




Xco:

5




10

Xcp:




15

Xcq:




20

Xcr:




25

Xcs:




-37-

Xct:




5

Xcu:



10

Xcv:



15

Xcw:


N-Methyl-N-ethyl-valine

Xcx:

N,N-Diethylvaline


20

Xcy:



25

Xcz:



-38-

Xda: N-Methyl-2-aminobutyroyl

Xdb: 2-aminobutyroyl


Xdc: N,N-Dimethyl-2-aminobutyroyl

5 Xdd: N,N-Dimethyl-2-tert.butylglycine


Xde: N,N-Dimethyl-isoleucine

Xdf: 2-tert.butylglycine

10 Xdg:




15 Xdh:




20

Xdi:



25


Xdk:



-39-


Xdl: N-Methyl-2-tert.butylglycine

Xdm:




5

Xdn:




10

Xdo:




15

Xdp:




20

Xdq:




25

Xdr:




Xds:




-40-


Xdt:




Xdu:




Xdv:




Xdw:




Xdx:



Xdy:




Xdz:






Xea:

5




Xeb:

10




Xec:

15




Xed:

20




Xee:

25




Xef:




-42-

Xeg:




5

Xeh:




10

Xei:




Xek:




15

Xel:




20

Xem:




25

Xen:




-43-

Xeo:




5

Xep:



10

Xeq:



15

Compounds of this invention may be assayed for anti-cancer activity by conventional methods, including for example, the methods described below.

20

#### A. In vitro methodology

Cytotoxicity was measured using a standard methodology for adherent cell lines such as the microculture tetrazolium assay (MTT). Details of this assay have been published (Alley, MC et al, Cancer Research 48:589-601, 1988).

25

Exponentially growing cultures of tumor cells such as the HT-29 colon carcinoma or LX-1 lung tumor are used to make microtiter plate cultures. Cells are seeded at 3000 cells per well in 96-well plates (in 150  $\mu$ l or media), and grown overnight at 37°C. Test compounds are added, in 10-fold dilutions

varying from  $10^{-4}$  M to  $10^{-10}$  M. Cells are then incubated for 72 hours. To determine the number of viable cells in each well, the MTT dye is added (50  $\mu$ l or 3 mg/ml solution of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide in saline). This mixture is incubated at 37°C for 5 hours, and then 50  $\mu$ l of 25% SDS, pH2 is added to each well. After an overnight incubation, the absorbance of each well at 550 nm is read using an ELISA reader. The values for the mean  $\pm$  SD of data from replicated wells are calculated, using the formula % T/C (% viable cells treated/control).

$$\frac{\text{OD of treated cells}}{\text{OD of control cells}} \times 100 = \% \text{ T/C}$$

The concentration of test compound which gives a T/C of 50% growth inhibition was designated as the  $IC_{50}$  value.

#### B. In vivo methodology

Compounds of this invention were further tested in pre-clinical assay for in vivo activity which is indicative of clinical utility. Such assays were conducted with nude mice into which tumor tissue, preferably of human origin, had been transplanted (xenografted), as is well known in this field. Test compounds were evaluated for their anti-tumor efficacy following administration to the xenograft-bearing mice.

More specifically, human breast tumors (MX-1) which had been grown in athymic nude mice were transplanted into new recipient mice, using tumor fragments which were about 50 mg in size. The day of transplantation was designated as day 0. Six to ten days later, mice were treated with the test compounds given as an intravenous injection or orally, in groups of 5-10 mice at

-45-

each dose. Compounds were given every other day, for 3 weeks, at doses from 1-200 mg/kg body weight.

- 5 Tumor diameters and body weights were measured twice weekly. Tumor volumes were calculated using the diameters measured with Vernier calipers, and the formula

$$(\text{Length} \times \text{width}^2)/2 = \text{mm}^3 \text{ of tumor volume}$$

- 10 Mean tumor volumes are calculated for each treatment group, and T/C values determined for each group relative to the untreated control tumors.

The new compounds possess good tumor inhibiting properties.